

**Expanded BRFSS, 2003**

**Oneida and Oswego Locality Tables**

TABLE OF CONTENTS:

Core 1: Health Status / Healthy Days .....	1
Core 2: Health Care Access .....	4
Core 3: Exercise .....	8
Core 4: Diabetes .....	9
Core 5: Asthma .....	10
Core 6: Arthritis .....	12
Core 7: Tobacco Use .....	14
Core 8: Tobacco ETS .....	17
Core 9: Alcohol Consumption .....	19
Core 10: Weight Status .....	21
Core 11: Mammography .....	23
Core 12: Sexual Behavior .....	25
Core 13: Family Planning .....	28
Core 14: Cardiovascular Disease .....	29
Core 15: Prostate Cancer Screening .....	31
Core 16: Colorectal Cancer Screening .....	37

**New York State Department Of Health**

Expanded BRFSS, 2003 - New York State Department of Health

## Self-rated Health Status

*Oneida, Oswego*

Demographic Groups	n <sup>1</sup>	Fair or Poor % <sup>2</sup>	n	Good to Excellent %	C.I. <sup>3</sup>
Total	91	11.6	559	88.4	2.6
Gender					
Male	29	10.4	217	89.6	3.9
Female	62	12.7	342	87.3	3.4
Age					
18-34	5	2.1	136	97.9	1.9
35-54	27	8.2	231	91.8	3.5
≥55	59	24.5	188	75.5	6.1
Education					
≤High School	60	16.5	249	83.5	4.4
>High School	31	7.0	308	93.0	2.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Reported Number of Physical Distress Days in the Past Month

*Oneida, Oswego*

Demographic Groups	n <sup>1</sup>	≥14 days % <sup>2</sup>	n	< 14 days %	C.I. <sup>3</sup>
Total	82	11.4	551	88.6	2.7
Gender					
Male	24	8.9	220	91.1	3.8
Female	58	13.8	331	86.2	3.9
Age					
18-34	9	6.1	132	93.9	4.5
35-54	29	8.6	226	91.4	3.5
≥55	44	20.2	189	79.8	5.9
Education					
≤High School	45	14.7	253	85.3	4.5
>High School	37	8.4	296	91.6	3.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Reported Number of Mental Distress Days in the Past Month

*Oneida, Oswego*

Demographic Groups	n <sup>1</sup>	≥14 days % <sup>2</sup>	n	< 14 days %	C.I. <sup>3</sup>
Total	69	10.2	570	89.8	2.6
Gender					
Male	16	6.4	229	93.6	3.3
Female	53	13.8	341	86.2	3.8
Age					
18-34	19	11.8	120	88.2	5.6
35-54	32	11.2	226	88.8	4.1
≥55	18	7.6	220	92.4	3.6
Education					
≤High School	40	13.4	261	86.6	4.4
>High School	29	7.2	307	92.8	2.8

<sup>1</sup> Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup> Weighted Percent

<sup>3</sup> 95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Health Insurance Coverage, 18-64 Years of Age

*Oneida, Oswego*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	431	84.6	66	15.4	3.8
Gender					
Male	161	80.9	35	19.1	6.2
Female	270	88.5	31	11.5	4.3
Age					
18-34	112	75.7	29	24.3	8.2
35-54	231	88.6	30	11.4	4.3
55-64	88	93.0	7	7.0	5.4
Education					
≤High School	173	76.6	45	23.4	6.6
>High School	258	91.5	21	8.5	4.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Health Insurance Coverage, Aged 18 and Older

*Oneida, Oswego*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	583	87.3	70	12.7	3.1
Gender					
Male	213	83.9	36	16.1	5.3
Female	370	90.6	34	9.4	3.4
Age					
18-34	112	75.7	29	24.3	8.2
35-54	231	88.6	30	11.4	4.3
≥55	236	96.2	11	3.8	2.4
Education					
≤High School	262	81.4	49	18.6	5.2
>High School	319	92.9	21	7.1	3.4

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Reported Having a Routine Physical Exam in Past Two Years

*Oneida, Oswego*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	569	86.6	80	13.4	3.1
Gender					
Male	203	82.1	45	17.9	5.2
Female	366	90.9	35	9.1	3.2
Age					
18-34	121	84.1	20	15.9	6.9
35-54	218	84.0	41	16.0	5.0
≥55	226	91.7	19	8.3	4.0
Education					
≤High School	264	83.9	45	16.1	4.8
>High School	303	89.1	35	10.9	3.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Time in Past 12 Months when Needed Medical Care but Could Not Get It

*Oneida, Oswego*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	39	5.5	613	94.5	2.0
Gender					
Male	11	4.3	238	95.7	3.1
Female	28	6.6	375	93.4	2.6
Age					
18-34	10	5.9	131	94.1	4.5
35-54	21	6.8	240	93.2	3.1
≥55	8	3.6	238	96.4	3.0
Education					
≤High School	18	6.3	292	93.7	3.4
>High School	21	4.7	319	95.3	2.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Participated in Leisure-Time Physical Activity or Exercise During the Past 30 Days

*Oneida, Oswego*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	487	77.3	165	22.7	3.5
Gender					
Male	201	81.7	48	18.3	5.3
Female	286	73.1	117	26.9	4.6
Age					
18-34	116	83.3	25	16.7	6.7
35-54	197	77.3	64	22.7	5.6
≥55	171	71.7	75	28.3	6.1
Education					
≤High School	211	70.3	99	29.7	5.7
>High School	275	84.1	65	15.9	4.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Physician Diagnosed Diabetes

*Oneida, Oswego*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	47	6.6	605	93.4	2.1
Gender					
Male	24	8.2	224	91.8	3.6
Female	23	5.1	381	94.9	2.3
Age					
18-34	0	0.0	141	100	0.0
35-54	12	4.2	249	95.8	2.8
≥55	35	15.7	211	84.3	5.5
Education					
≤High School	27	8.3	283	91.7	3.5
>High School	19	4.8	321	95.2	2.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Lifetime Asthma Diagnosed by a Medical Professional

*Oneida, Oswego*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	75	12.0	576	88.0	3.0
Gender					
Male	17	8.5	231	91.5	4.1
Female	58	15.5	345	84.5	4.2
Age					
18-34	20	15.9	121	84.1	7.0
35-54	33	11.8	228	88.2	4.4
≥55	21	8.3	224	91.7	3.7
Education					
≤High School	39	13.5	270	86.5	4.7
>High School	35	10.5	305	89.5	3.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Current Asthma Diagnosed by a Medical Professional

*Oneida, Oswego*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	55	8.1	594	91.9	2.4
Gender					
Male	8	3.6	239	96.4	2.7
Female	47	12.5	355	87.5	3.8
Age					
18-34	11	7.8	129	92.2	4.9
35-54	25	8.8	236	91.2	3.9
≥55	18	7.2	226	92.8	3.5
Education					
≤High School	28	9.3	279	90.7	3.9
>High School	26	7.0	314	93.0	2.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Arthritis Diagnosed by a Medical Professional

*Oneida, Oswego*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	236	32.8	415	67.2	4.0
Gender					
Male	74	28.3	174	71.7	6.1
Female	162	37.1	241	62.9	5.1
Age					
18-34	16	8.9	125	91.1	4.7
35-54	82	31.3	178	68.7	6.3
≥55	138	57.2	108	42.8	6.9
Education					
≤High School	129	36.7	181	63.3	6.0
>High School	106	29.1	233	70.9	5.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Adults with Doctor Diagnosed Arthritis or Possible Arthritis (Chronic Joint Symptoms Only)

*Oneida, Oswego*

Demographic Groups	n <sup>1</sup>	Doctor Diagnosed % <sup>2</sup>	C.I. <sup>3</sup>	n	Possible Arthritis (CJS Only) %	C.I.	n	No Arthritis or CJS %	C.I.
Total	236	32.8	4.0	94	14.6	3.1	320	52.6	4.4
Gender									
Male	74	28.3	6.1	41	15.7	4.9	133	56.0	6.8
Female	162	37.2	5.1	53	13.5	3.7	187	49.3	5.4
Age									
18-34	16	8.9	4.7	24	17.7	7.0	101	73.4	7.9
35-54	82	31.3	6.3	50	18.9	5.2	128	49.7	6.8
≥55	138	57.4	6.9	19	6.5	3.1	88	36.1	6.7
Education									
≤High School	129	36.7	6.0	51	16.7	4.8	130	46.6	6.4
>High School	106	29.1	5.4	43	12.7	3.9	189	58.2	5.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Ever Smoked 100 Cigarettes in Lifetime

*Oneida, Oswego*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	353	54.7	299	45.3	4.4
Gender					
Male	156	60.7	93	39.3	6.8
Female	197	48.9	206	51.1	5.4
Age					
18-34	83	59.9	58	40.1	9.1
35-54	139	52.0	121	48.0	6.8
≥55	129	53.1	118	46.9	7.0
Education					
≤High School	179	58.7	131	41.3	6.3
>High School	172	50.8	168	49.2	6.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Current Smoker

*Oneida, Oswego*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	172	27.9	480	72.1	4.0
Gender					
Male	74	31.7	175	68.3	6.6
Female	98	24.2	305	75.8	4.6
Age					
18-34	58	41.5	83	58.5	9.2
35-54	79	29.1	181	70.9	6.1
≥55	34	14.1	213	85.9	4.8
Education					
≤High School	101	36.0	209	64.0	6.2
>High School	71	20.4	269	79.6	5.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Status of Cigarette Smoking***Oneida, Oswego*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Everyday Smoker %<sup>2</sup></b>	<b>C.I.<sup>3</sup></b>	<b>n</b>	<b>Someday Smoker %</b>	<b>C.I.</b>	<b>n</b>	<b>Former or Never Smoked %</b>	<b>C.I.</b>
Total	131	20.2	3.5	41	7.7	2.6	480	72.1	4.0
Gender									
Male	52	20.9	5.6	22	10.8	4.8	175	68.3	6.6
Female	79	19.5	4.3	19	4.7	2.4	305	75.8	4.6
Age									
18-34	37	24.2	7.7	21	17.3	7.5	83	58.5	9.2
35-54	67	24.6	5.8	12	4.5	2.8	181	70.9	6.1
≥55	26	11.2	4.4	8	2.8	2.2	213	85.9	4.8
Education									
≤High School	85	29.4	5.8	16	6.6	3.4	209	64.0	6.2
>High School	46	11.6	3.6	25	8.8	4.0	269	79.6	5.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution<sup>2</sup>Weighted Percent<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Home Smoking Policies**

*Oneida, Oswego*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Not Allowed %<sup>2</sup></b>	<b>C.I.<sup>3</sup></b>	<b>n</b>	<b>Allowed Some Places %</b>	<b>C.I.</b>	<b>n</b>	<b>Allowed Anywhere %</b>	<b>C.I.</b>
Total	387	61.4	4.2	93	14.1	3.1	171	24.5	3.7
Gender									
Male	141	58.8	6.8	31	13.8	5.0	76	27.4	6.0
Female	246	63.9	5.1	62	14.4	3.6	95	21.7	4.3
Age									
18-34	86	61.2	9.0	24	16.2	6.9	31	22.5	7.6
35-54	162	64.8	6.4	38	12.3	4.1	60	22.9	5.7
≥55	137	57.9	6.9	30	14.1	5.3	79	28.0	6.0
Education									
≤High School	167	54.2	6.4	46	16.7	5.0	96	29.1	5.7
>High School	219	68.3	5.5	46	11.5	3.7	75	20.2	4.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Home Smoking

*Oneida, Oswego*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	188	29.7	465	70.3	4.1
Gender					
Male	78	33.4	171	66.6	6.7
Female	110	26.1	294	73.9	4.6
Age					
18-34	47	36.0	94	64.0	9.2
35-54	75	27.6	186	72.4	6.0
≥55	63	25.5	184	74.5	6.1
Education					
≤High School	111	37.3	200	62.7	6.2
>High School	77	22.6	263	77.4	5.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**At Risk for Binge Drinking (Drank in Past Month and had 5 or more Drinks per drink Occasion)**

*Oneida, Oswego*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	92	17.1	553	82.9	3.6
Gender					
Male	61	25.9	182	74.1	6.3
Female	31	8.8	371	91.2	3.4
Age					
18-34	28	22.2	113	77.8	8.1
35-54	49	22.2	207	77.8	6.0
≥55	15	6.7	229	93.3	3.6
Education					
≤High School	49	20.0	258	80.0	5.4
>High School	43	14.4	293	85.6	4.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**At Risk for Heavy Drinking (Males More Than 2 Drinks and Females More Than 1 Drink per Day in Past Month)**

*Oneida, Oswego*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	35	6.5	612	93.5	2.5
Gender					
Male	20	9.1	225	90.9	4.4
Female	15	4.0	387	96.0	2.3
Age					
18-34	11	10.8	128	89.2	6.7
35-54	14	5.1	245	94.9	2.8
≥55	10	4.4	235	95.6	3.0
Education					
≤High School	14	5.7	294	94.3	3.1
>High School	21	7.3	316	92.7	3.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Overweight or Obese, BMI(Body Mass Index) 25 or More**

*Oneida, Oswego*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	384	61.7	244	38.3	4.4
Gender					
Male	179	71.9	67	28.1	6.4
Female	205	51.6	177	48.4	5.6
Age					
18-34	67	48.3	69	51.7	9.5
35-54	159	67.1	89	32.9	6.4
≥55	157	68.4	84	31.6	6.4
Education					
≤High School	189	60.9	114	39.1	6.4
>High School	194	62.6	129	37.4	6.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Weight Status (Not Overweight BMI<25, Overweight BMI 25-29, Obese BMI 30+)**

*Oneida, Oswego*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Underweight\ Normal %<sup>2</sup></b>	<b>C.I.<sup>3</sup></b>	<b>n</b>	<b>Overweight %</b>	<b>C.I.</b>	<b>n</b>	<b>Obese %</b>	<b>C.I.</b>
Total	244	38.3	4.4	233	36.0	4.2	151	25.7	4.0
Gender									
Male	67	28.1	6.4	109	41.6	6.8	70	30.3	6.5
Female	177	48.4	5.6	124	30.5	5.0	81	21.1	4.5
Age									
18-34	69	51.7	9.5	42	28.0	8.2	25	20.3	7.8
35-54	89	32.9	6.4	89	35.9	6.7	70	31.1	6.6
≥55	84	31.6	6.4	101	43.7	7.0	56	24.7	6.3
Education									
≤High School	114	39.1	6.4	119	37.5	6.1	70	23.4	5.5
>High School	129	37.4	6.0	113	34.5	5.8	81	28.1	5.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Ever Had Mammography Screening, Women Aged 40 Years and Older**

*Oneida, Oswego*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Female	248	89.5	32	10.5	3.7
Age					
40-64	157	88.5	24	11.5	4.7
≥65	91	91.5	8	8.5	6.1
Education					
≤High School	118	89.7	14	10.3	5.5
>High School	129	89.2	18	10.8	5.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Received Mammography Screening in Past Two Years, Women Aged 40 Years and Older**

*Oneida, Oswego*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Female	214	77.0	65	23.0	5.4
Age					
40-64	138	76.8	43	23.2	6.7
≥65	76	77.5	22	22.5	9.0
Education					
≤High School	102	76.7	30	23.3	7.9
>High School	112	77.6	34	22.4	7.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Ever Counseled by Medical Professional on Prevention of STD's Through Condom Use, Aged 64 or Younger**

***Oneida, Oswego***

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	74	16.2	416	83.8	4.0
Gender					
Male	20	12.0	174	88.0	5.6
Female	54	20.6	242	79.4	5.5
Age					
18-34	44	30.6	95	69.4	8.7
35-54	23	8.7	235	91.3	3.8
55-64	7	5.1	86	94.9	4.1
Education					
≤High School	41	20.0	174	80.0	6.2
>High School	33	12.9	242	87.1	5.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Sexually Active, Age 18-64 Years Old**

*Oneida, Oswego*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	360	83.5	91	16.5	3.9
Gender					
Male	146	83.2	33	16.8	6.2
Female	214	83.8	58	16.2	4.6
Age					
18-34	117	85.8	15	14.2	7.4
35-54	195	86.4	41	13.6	4.6
55-64	48	68.5	35	31.5	10.6
Education					
≤High School	156	84.4	41	15.6	5.3
>High School	204	82.7	50	17.3	5.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Used Condom Last Time Had Sexual Intercourse, Aged 64 or Younger

*Oneida, Oswego*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	72	18.6	295	81.4	4.6
Gender					
Male	37	21.1	115	78.9	7.1
Female	35	15.8	180	84.2	5.7
Age					
18-34	33	27.2	87	72.8	9.3
35-54	35	14.7	163	85.3	5.1
55-64	4	7.1	45	92.9	7.1
Education					
≤High School	37	21.3	125	78.7	7.1
>High School	35	16.1	170	83.9	5.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Birth Control(BC) Use To Prevent Pregnancy Among Reproductive Aged Men(18-59) and Women(18-44)

*Oneida, Oswego*

Demographic Groups	Use BC			No BC at risk			No BC		
	n <sup>1</sup>	% <sup>2</sup>	C.I. <sup>3</sup>	n	%	C.I.	n	%	C.I.
Total	197	58.8	6.2	44	14.8	4.5	81	26.4	5.7
Gender									
Male	93	52.3	8.5	24	16.3	6.4	49	31.4	8.1
Female	104	69.2	8.0	20	12.6	5.8	32	18.3	6.6
Age									
18-34	81	56.2	9.6	25	19.2	7.5	30	24.6	8.9
35-59	116	61.3	7.9	19	10.7	5.2	51	28.0	7.3
Education									
≤High School	86	59.4	9.1	19	16.1	7.4	36	24.5	7.8
>High School	111	58.2	8.4	25	13.7	5.5	45	28.0	8.2

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Physician Diagnosed Heart Attack, Angina or Stroke**

*Oneida, Oswego*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	59	8.2	588	91.8	2.3
Gender					
Male	30	10.6	217	89.4	4.0
Female	29	5.9	371	94.1	2.3
Age					
18-64	26	4.5	467	95.5	1.9
≥65	33	23.8	118	76.2	7.8
Education					
≤High School	31	8.4	276	91.6	3.2
>High School	28	7.9	310	92.1	3.2

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Physician Diagnosed Coronary Heart Disease (Heart Attack or Angina)**

*Oneida, Oswego*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	43	6.1	603	93.9	2.0
Gender					
Male	26	8.7	221	91.3	3.6
Female	17	3.6	382	96.4	1.9
Age					
18-64	21	3.6	472	96.4	1.7
≥65	22	16.6	128	83.4	7.0
Education					
≤High School	22	6.5	284	93.5	3.0
>High School	21	5.7	317	94.3	2.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Ever Had Prostate Specific Antigen Test, Men Aged 40 and Older

*Oneida, Oswego*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Male	86	53.5	72	46.5	8.5
Age					
40-64	49	44.6	62	55.4	10.0
≥65	37	79.8	10	20.2	12.2
Education					
≤High School	42	46.6	37	53.4	12.1
>High School	43	60.2	35	39.8	11.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Received Prostate Specific Antigen Test in the Past Two Years, Men Aged 40 and Older

*Oneida, Oswego*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Male	71	45.7	85	54.3	8.5
Age					
40-64	37	35.4	73	64.6	9.8
≥65	34	76.4	12	23.6	12.9
Education					
≤High School	35	41.4	43	58.6	11.9
>High School	35	49.6	42	50.4	12.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Ever Had a Digital Rectal Exam, Men Aged 40 and Older

*Oneida, Oswego*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Male	129	76.0	39	24.0	7.3
Age					
40-64	88	74.1	29	25.9	9.0
≥65	41	81.8	10	18.2	11.2
Education					
≤High School	60	70.6	24	29.4	11.2
>High School	68	81.2	15	18.8	9.4

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Ever Used a Blood Stool Test at Home, Aged 50 and Older**

*Oneida, Oswego*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	147	50.4	149	49.6	6.3
Gender					
Male	42	43.1	61	56.9	10.5
Female	105	56.3	88	43.7	7.6
Age					
50-64	77	52.1	72	47.9	9.0
≥65	70	48.5	77	51.5	8.9
Education					
≤High School	77	49.9	81	50.1	8.7
>High School	70	51.6	66	48.4	9.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Used a Blood Stool Test at Home in the Past Year, Aged 50 and Older**

*Oneida, Oswego*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	59	20.7	234	79.3	5.3
Gender					
Male	16	16.4	86	83.6	7.9
Female	43	24.1	148	75.9	7.0
Age					
50-64	32	20.8	116	79.2	7.4
≥65	27	20.5	118	79.5	7.5
Education					
≤High School	30	20.2	126	79.8	7.1
>High School	29	21.5	106	78.5	7.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Used a Blood Stool Test at Home in the Past 2 Years, Aged 50 and Older

*Oneida, Oswego*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	87	30.7	206	69.3	6.0
Gender					
Male	26	27.5	76	72.5	9.6
Female	61	33.3	130	66.7	7.5
Age					
50-64	49	31.8	99	68.2	8.3
≥65	38	29.6	107	70.4	8.6
Education					
≤High School	46	30.4	110	69.6	8.1
>High School	41	31.4	94	68.6	8.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Ever Had a Sigmoidoscopy or Colonoscopy, Aged 50 and Older

*Oneida, Oswego*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	145	49.6	151	50.4	6.3
Gender					
Male	48	47.4	54	52.6	10.5
Female	97	51.3	97	48.7	7.7
Age					
50-64	63	43.7	86	56.3	8.9
≥65	82	56.3	65	43.7	8.8
Education					
≤High School	75	46.6	82	53.4	8.7
>High School	69	52.6	68	47.4	9.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Had a Sigmoidoscopy or Colonoscopy in the Past 10 Years, Aged 50 and Older

*Oneida, Oswego*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	135	45.6	159	54.4	6.3
Gender					
Male	45	44.1	56	55.9	10.5
Female	90	46.8	103	53.2	7.7
Age					
50-64	57	39.1	91	60.9	8.8
≥65	78	53.1	68	46.9	8.9
Education					
≤High School	69	42.8	86	57.2	8.6
>High School	65	48.4	72	51.6	9.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Had a Blood Stool Test Using a Home Kit in the Past Year or Had a Lower Endoscopy in the Past 10 Years, Aged 50 and Older**

***Oneida, Oswego***

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	156	52.3	143	47.7	6.3
Gender					
Male	48	47.2	56	52.8	10.5
Female	108	56.4	87	43.6	7.6
Age					
50-64	68	45.7	81	54.3	9.0
≥65	88	59.8	62	40.2	8.6
Education					
≤High School	80	48.7	80	51.3	8.7
>High School	75	56.2	62	43.8	9.2

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval